State of Alaska Department of Fish and Game Homination for Waters Important to Anadromous Fish

	alog Number of Waterway	. 7 7 3 1	nece	me - Tor	al name
e of Waterway					
lition X Delet	ion Correction _	Backup	Informacio	"	
	for the state of t	Office Use		. 1	
Nomination # 94 316 Revision Year:			LandShea		
		Regional Supervisor			Date
Revision Year:	sCatalog		1.2-		11/19/93
Revision to: Atla			200	77	1-1-
Revision Code: A - Z		2. Drove			
Revision Code:		Drafted			Date
T PER MILLE	OPCERVAT	ION INFORMAT	TON		
	Date(s) Observed		Rearing	Migration	Anadromous
Species CoHo	7-23-93		4 *		YES
CUTTHROAT	" 1		>10		NO
COLLINGAL					
PORTANT: Provide a	all supporting document migration of anadromo	tation that us fish, inc	this water luding: num	body is import of fish a	ortant for the
portant: Provide a pawning, rearing or oserved; sampling m stach a copy of a m swell as any other paring habitat; locomments: BEAVEL DAM	migration of anadromomethods, sampling durated ap showing location of information such as: cations, types, and he was STROAM /LAKE SYSTEM	ion and area mouth and o specific st ights of any	sampled; observed upgiveam reach barriers;	copies of fie per extent of es observed etc.	each species, as spawning or
portant: Provide a bawning, rearing or oserved; sampling m stach a copy of a m swell as any othe earing habitat; loc omments: BEAVER DAM	migration of anadromomethods, sampling durated ap showing location of information such as: cations, types, and he was STREAM /LAKE SYSTEMED STREAM /LAKE SYSTEMED STREAM /LAKE SYSTEMED AT SMITT OUTLET AT SMITTERS	ion and area mouth and o specific stights of any m. ANAUROM	sampled; observed up; ream reach barriers; ous HABITAT	copies of fie per extent of es observed etc. EXTENDS APPRO BEAVER DAM	each species, as spawning or KIMATEY (APPROX. 5'
portant: Provide a sawning, rearing or served; sampling match a copy of a match as any other sampling habitat; local match as a sample of the	migration of anadromomethods, sampling durates ap showing location of rinformation such as: cations, types, and he med STREAM /LAKE SYSTEMED STREAM /LAKE SYSTEMED STREAM /LAKE SYSTEMED AT SMITTER AT	ion and area mouth and o specific stights of any om. ANADROMATH LAGOON TO BAITED MINN	sampled; of beerved upported to the harriers; out the high of the	copies of fie ber extent of es observed etc. EXTENDS APPRO BEAVER DAM	Id notes; etc. each species, as spawning or XIMATEY (APPROX. 5'
portant: Provide a pawning, rearing or pserved; sampling metach a copy of a a	migration of anadromomethods, sampling durated ap showing location of rinformation such as: cations, types, and he med stream /LAKE SYSTE FROM ITS OUTLET AT SMIRE MEAN MIGRATION). THERE A COLLECTIVEY YIELDES A	ion and area mouth and o specific st ights of any m. ANADROM THE BAITED MINN TOTAL OF 4	sampled; observed up; ream reach barriers; ous HABITAT of THE FIRST ow TRAAS SE COHO FRY,	es observed etc. EXTENDS APPROPRIET AT SUCCESS NUMEROUS CA	Id notes; etc. each species, as spawning or EXIMATELY (APPROX. 5' GIVE LOCATIONS OFTHROAT TROUT
portant: Provide a sawning, rearing or served; sampling metach a copy of a metach a sampling habitat; locomments: BEAVER DAME, 000 FEET UPSTREAM HEH BARRIER TO UPSTREAM OF DAME, DUNSTREAM OF DAME, DUNSTREAM OF DAME, NO 1 SCULPIN, TWO	migration of anadromomethods, sampling durates ap showing location of information such as: cations, types, and he med stream /LAKE SYSTE FROM ITS OUTLET AT SMIREAM MIGRATION). THERE A COLLECTIVEY YIELDES ATTRAPS SET WITHIN THE LA	ion and area mouth and o specific strights of any om. ANADROM THE BAITED MINN TOTAL OF 4 KE ABOVE THE	sampled; cobserved upgream reach barriers; ous HABITAT of THE FIRST ow TRAPS SE COHO FRY, E BEAVER DA	expies of fie ber extent of es observed etc. EXTENDS APPRO- BEAVER DAM IT AT SUCCESS NUMEROUS CO M YIELDED A S	Id notes; etc. each species, as spawning or eximately (APPROX. 5' LIVE LOCATIONS OFTHROAT TROUT
portant: Provide a waring, rearing or tach a copy of a maring habitat; localing habi	migration of anadromomethods, sampling duration per showing location of rinformation such as: cations, types, and he med stream /LAKE SYSTE FROM ITS OUTLET AT SMIRE AMORATION). THERE A COLLECTIVEY YIELDED A TRAPS SET WITHIN THE LA ATTACHED JULY 27, 1992	ion and area mouth and o specific stagets of any om. ANADROMATH LAGOON TO BAITED MINN TOTAL OF 4 KE ABOVE THE BEST MACLOCATIONS LIMING AND ANADROMATICAL OF THE BEST MACLOCATIONS LIMING AND	sampled; cobserved upgream reach barriers; ous HABITAT of the First ow traps se coho free, beaver Date to the first of the first own traps se coho free, beaver Date to the first to the fi	copies of fie ber extent of es observed etc. EXTENDS APPRO- BEAVER DAM TAT SUCCESS NUMEROUS CO M YIELDED A S TON REPORT.	Id notes; etc. each species, as spawning or eximately (APPROX. 5' LIVE LOCATIONS OFTHROAT TROUT
portant: Provide a awning, rearing or served; sampling match a copy of a match as any other well as any other aring habitat; local match as a served. Daments: Beaver Daments:	migration of anadromomethods, sampling duration per showing location of rinformation such as: cations, types, and he med stream /LAKE SYSTE FROM ITS OUTLET AT SMIRE AMORATION). THERE A COLLECTIVEY YIELDED A TRAPS SET WITHIN THE LA ATTACHED JULY 27, 1992	ion and area mouth and o specific stagets of any om. ANADROMATH LAGOON TO BAITED MINN TOTAL OF 4 KE ABOVE THE BEST MACLOCATIONS LIMING AND ANADROMATICAL OF THE BEST MACLOCATIONS LIMING AND	sampled; cobserved upgream reach barriers; ous HABITAT of the First ow traps se coho free, beaver Date to the first of the first own traps se coho free, beaver Date to the first to the fi	copies of fie ber extent of es observed etc. EXTENDS APPRO- BEAVER DAM TAT SUCCESS NUMEROUS CO M YIELDED A S TON REPORT.	Id notes; etc. each species, as spawning or eximately (APPROX. 5' LIVE LOCATIONS OFTHROAT TROUT
PORTANT: Provide a awning, rearing or served; sampling match a copy of a a cop	migration of anadromomethods, sampling durates ap showing location of rinformation such as: cations, types, and he med STROAM /LAKE SYSTEMED STROAM /LAKE SYSTEMED STROAM /LAKE SYSTEMED AND COLLECTIVELY YIELDED ATTRAPS SET WITHOUT THE LATTACHED JULY 27, 1992 CTREMEN LOW ~ TRAPPING COLLECTIVELY LOW ~ TRAPPING COLLECTIVELY YIELDED ATTRAPS SET WITHOUT THE LATTACHED JULY 27, 1992 CTREMEN LOW ~ TRAPPING COLLEGE PRINTS (KEVIN) J	ion and area mouth and o specific stagets of any om. ANADROMATH LAGOON TO BAITED MINN TOTAL OF 4 KE ABOVE THE BEST MACLOCATIONS LIMING AND ANADROMATICAL OF THE BEST MACLOCATIONS LIMING AND	sampled; cobserved upgream reach barriers; ous HABITAT of the First ow traps se coho free, beaver Date to the first of the first own traps se coho free, beaver Date to the first to the fi	copies of fie ber extent of es observed etc. EXTENDS APPRO- BEAVER DAM THE SUCCESS NUMEROUS CA MYIBLDED A SE TOWN REPORT. W REMAINANTS HOLOGIST	APPROX. 5' CAPPROX. 5' CAPPROX. 5' CAPPROX. 5' CAPPROX. TRAVE CAPPROX. TR
PORTANT: Provide a awning, rearing or served; sampling match a copy of a a cop	migration of anadromomethods, sampling durates ap showing location of information such as: cations, types, and he med STREAM /LAKE SYSTEMED STREAM MICRATION). THREE COLLECTIVEY YIELDES ATTRAPS SET WITHIN THE LA ATTACHED JULY 27, 1992 (TREMEN LOW ~ TRAPPING OF TRAPPING O	ion and area mouth and o specific stages of any om. ANADROMUTH LAGOON TO BAITED MINN TOTAL OF 4 KE ABOVE THE BEST MACLOCATIONS LIMITED THANLEY, A. Hanley	I SAMPLED; Cobserved upgream reach barriers; ous HABITAT O THE FIRST OW TRAPS SE COHO FRY, E BEAVER DATES INSPECTED TO THE FE HABITAT B	copies of fie ber extent of es observed etc. EXTENDS APPRO- BEAVER DAM IT AT SUCCESS NUMEROUS CO M YIELDED A S MON REPORT. EN REMANNIES IOLOGIST	Id notes; etc. each species, as spawning or EXIMATELY (APPROX. 5' EVE LOCATIONS OTTHROAT TROUT ENGLE JOILY FOOLS ALASKA DEPT. FISH & GAM 100 0 5 19
portant: Provide a waring, rearing or tach a copy of a maring habitat; locaring habitation of DAm habitation of DAm habitation characteristics.	migration of anadromomethods, sampling durates ap showing location of information such as: cations, types, and he med STREAM /LAKE SYSTEMED STREAM MICRATION). THREE COLLECTIVEY YIELDES ATTRAPS SET WITHIN THE LA ATTACHED JULY 27, 1992 (TREMEN LOW ~ TRAPPING OF TRAPPING O	ion and area mouth and o specific stages of any om. Anadrom the Lacon to the Baited minn total of 4 Ke above the Best Irac catrons limit. Hanley	I SAMPLED; Cobserved upgream reach barriers; ous HABITAT O THE FIRST OW TRAPS SE COHO FRY, E BEAVER DATES INSPECTED TO THE FE HABITAT B	EXTENDS APPLE EXTENDS APPLE BEAVER DAM IT AT SUCCESS NUMEROUS CA MYIBLED A S TON REPORT. WILLIAM LOLOGIST DIVISION,	APPROX. 5' CAPPROX. 5' CAPPROX. 5' CAPPROX. 5' CAPPROX. TRAVE CAPPROX. TR

MEMORANDUM

STATE OF ALASKA DEPARTMENT OF FISH AND GAME

To: Al Peterson

DATE: July 27, 1993

Forest Practices Forester

Department of Natural FILE No: SE-88-001

Resources Ketchikan

PHONE: 225-2027

FROM:

Kevin J. Hanley KgH SUBJECT: Forest Practices
Habitat Biologist Inspection -

Habitat Biologist Inspection Habitat and Restoration Division ITT Rayonier Smith

Ketchikan Cove

On July 23, 1993, you, Clarence Clark, and I conducted a Forest Practices Inspection at ITT Rayonier's Smith Cove tract to examine a series of streams and lakes within or adjacent to Units 93-9.1A, 93-9.1B, 93-10.1, and 93-11.1 for the purpose of determining their waterbody classifications and riparian protection needs. The results of this inspection are as follows:

We began by examining the reach of stream located between the southern boundary of Unit 93-10.1 and the northern boundary of Unit 93-11.1. This reach is the lowermost portion of the stream/lake system that flows through and adjacent to the eastern portion of Unit 93-10.1. With the exception of a bedrock controlled portion of the south bank in the vicinity of the flagged "F" Spur alignment crossing, the stream banks are predominantly vegetatively controlled and the substrate ranges from organic material to large and small angular cobbles with lesser amounts of gravel. Stream gradient averages less than 8 percent throughout, ranging from 2 percent in the area just downstream of the lake outlet, to 5 percent in the vicinity of the "F" Spur alignment crossing. The upper reach of this system between the 2000 Mainline Road and the upper lake was previously trapped on June 22, 1993 and determined to be nonanadromous by the capture of only resident cutthroat trout, Dolly Varden char, and three-spine stickleback. It was assumed at that time that the beaver dammed outlets of the two lakes effectively precluded anadromous fish from accessing and utilizing upstream habitat. This assumption was verified by the results of the trapping that was conducted during this most recent inspection.

A total of three baited minnow traps were set at successive downstream locations within this lower reach between the beaver dammed outlet of the lower lake and the stream's saltwater mouth at Smith Lagoon. Collectively, these traps yielded a total of four coho juveniles and numerous cutthroat trout. Two additional traps were set within the lake near its beaver dammed outlet. Only one of these yielded a single

Dolly Varden char. As such, The 5-foot high, nearly vertical beaver dam can be assumed to be a barrier to anadromous fish. Blue/white "Watercourse and Lake Protection Zone" ribbon was hung throughout the Type A habitat downstream of the dam to identify the waters along which 66-foot no-cut riparian buffers are required. In addition to the mainstem, these include the off-channel habitat located just upstream of the "F" Spur alignment crossing and the low gradient tributary which enters the main channel from the shallow beaver dammed lake located on the east side of Unit 93-11.1. The large, well vegetated dam at this lake's outlet appears to form an effective barrier to the upstream migration of coho fry and, therefore, defines the upper extent of type A classification within the tributary.

According to the notification map, a 48-inch CMP culvert is proposed for the "F" Spur alignment crossing of the main channel. Although not discussed during the inspection, this culvert must be bedded at a sufficient depth (10") below the streambed to prevent perching and provide for the efficient passage of fish throughout.

On a separate note, two of the three traps that were set within this lower type A reach were located within the northeastern portion of clearcut Unit 133, which was notified on June 11, 1991 and apparently harvested soon after without retaining a buffer. The notification materials submitted for this unit gave no indication as to the existence, location and classification of this 5 to 10 foot wide type A stream. As such, the state was not aware of its existence until the time of this inspection, approximately two years after it was clearcut to the banks. Like the upper reach of Hole-in-the-Wall Creek at Lyman Anchorage, this failure to notify the location and classification of known surface waters has resulted in the degradation of fish habitat. Although it is difficult to determine the effects of this bank-to-bank clearcutting in the relatively short period of time since it occurred, it can be assumed that the lack of shade has substantially increased stream temperatures. This is especially critical to the survival of fish during low stream flow conditions such as those that we have been experiencing this summer. The finding of a dead rough skinned newt within one of the pools in which coho fry were trapped may be indicative of the effects of elevated stream temperatures. Although nothing can be done to mitigate the loss of this riparian zone, ITT Rayonier apparently is now aware of the necessity for locating and classifying all surface waters in their areas of operation prior to the commencement of harvest activities.

In addition to the bank-to-bank clearcutting of Unit 133, past harvesting of the unit located on the east side of the lower lake deposited substantial amounts of trees into both the lake and the lower type A reach in the vicinity of the beaver dam.

Al Peterson 3 July 27, 1993

Although this harvesting occurred prior to the 1990 revision of the Forest Practices Act, it is a prime example of poor management practices and non-adherence to BMP's. We trust that this type of activity will not occur in the future.

Following our examination of this lower reach, we proceeded to the lake located in the south-central portion of Unit 93-9.1A. Two baited minnow traps were set beneath the scant amount of large woody debris cover near the lake's beaver dammed outlet. After an approximate 1 hour soak, both traps failed to yield fish of any kind. As such, this lake can be assumed to be a nonanadromous waterbody.

We closed out this inspection by examining the two lakes located within and adjacent to Unit 93-9.1B. Time constraints and an exhausted supply of bait prevented the trapping of both lakes. However, one baited minnow trap was set within the lower lake and allowed to soak for approximately 20 minutes. This trap yielded no fish and, therefore, both lakes can be assumed to be nonanadromous. Incidentally, the short reach of stream connecting these two lakes contains substantial amounts of trees that were felled into and across the stream during the selective helicopter harvesting of Unit 171. This unit was notified on March 19, 1991 and, as with clearcut Unit 133, the notification materials failed to notify the existence, location and classification of this stream. The deposition of trees into this stream is especially objectionable given the fact that the unit was selectively harvested and, as such, greater opportunity should have existed to either retain this non-merchantable timber or directionally fall it away from the stream. We would request that it be removed from the stream during the harvest of Unit 93-9.1B.

Although the lakes examined during this inspection are assumed to be nonanadromous, they all harbor populations of resident fish. In addition, they may provide important habitat for Vancouver Canada geese and other waterfowl. As such, we would like to request and recommend that buffers of at least low-value and non-merchantable trees be retained along their shorelines and that commercial timber be directionally felled into the units. These practices would help to ensure the maintenance of water quality and the protection of resident fish habitat and would also provide a semblance of residual nearshore upland habitat.

Thanks for arranging this inspection. If you have any need to discuss this report, please contact me at 225-2027.

cc: L. Shea, ADF&G, Douglas

J. Gustafson, ADF&G, Ketchikan

J. Ferguson, ADEC, Juneau

C. Clark, ITT Rayonier, Inc., Ketchikan

P. Valentine, Zeman Logging N., Ketchikan C. Doig, Forest Management, Inc., Olympia, WA

L. Thompson, Kavilco, Inc., Kasaan





